



What if cell culture media do not mimic *in vivo* redox settings?

Ivan Spasojević

To cite this article: Ivan Spasojević (2016) What if cell culture media do not mimic *in vivo* redox settings?, Redox Report, 21:3, 127-129, DOI: [10.1179/1351000215Y.0000000036](https://doi.org/10.1179/1351000215Y.0000000036)

To link to this article: <https://doi.org/10.1179/1351000215Y.0000000036>



Published online: 18 Feb 2016.



Submit your article to this journal [↗](#)



Article views: 1512



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

What if cell culture media do not mimic *in vivo* redox settings?

Ivan Spasojević

Department of Life Sciences, Institute for Multidisciplinary Research, University of Belgrade, Belgrade, Kneza Višeslava 1, Serbia

Here, I address the topic of suitability for redox research of common settings in cell cultures. This is done through the prism of *in vitro* anticancer effects of vitamin C. Cell culture media show lower concentrations of iron and a higher level of oxygen compared to interstitial fluid. Such a setup promotes ascorbate-mediated production and accumulation of hydrogen peroxide, which efficiently kills a variety of cancer cell lines. However, the anticancer effects are annihilated if the iron level is corrected to mimic *in vivo* concentrations. It appears that the potential benefits of application of vitamin C in cancer treatment have been significantly overestimated. This might be true for other pro-oxidative agents as well, such as some (poly)phenols. We urgently need to establish medium formula and culture maintenance settings that are optimal for redox research.

Keywords: Cell culture, Iron, Ascorbate, Cancer

Commercial cell culture media, such as DMEM or RPMI-1640 supplemented with (usually) 10% (v/v) fetal calf serum (FCS), are built to mimic extracellular milieu, i.e. interstitial fluid. Apparently, on many accounts they do. Different types of cultured cells efficaciously grow and proliferate in those media, and there are thousands of cell culture studies conducted each year, some of them in the redox field. For example, about 250 papers were published in 2014, containing a combination of terms ‘antioxidant’ in the Title/Abstract section and ‘cell culture’ in All Fields (source: PubMed). Pertinent to this, it is clearly of the essence that cell culture media reflect *in vivo* redox settings as accurately as possible. But what if they do not, and what might be the consequences?

It is believed that the concentration of iron in interstitial fluid (outside the CNS) closely mirrors the range (10–30 μM) found in the plasma.^{1,2} The presence of different complexes and the susceptibility of iron in interstitial fluid to reduction and oxidation are not known, but cell culture media fail to reproduce even the concentration. DMEM contains only 0.25 μM ferric nitrate, whereas in RPMI-1640 iron probably exists via impurities. The level of iron in fetal calf serum (FCS) is not consistent and varies between manufacturers and batches. In the end, the concentration of iron does not exceed 5 μM in cell culture media

with 10% (v/v) FCS.^{3,4} The concentration of oxygen represents another major problem of redox research in cell cultures, as repeatedly pointed out by Halliwell.^{5,6} Cell cultures are kept under 95/5% air/CO₂ atmosphere resulting in hyperoxia of the medium ($\text{pO}_2 \approx 150$ mmHg) as compared to interstitial fluid, where pO_2 is in the 1–10 mmHg range.

Following over 20 studies on cancer cell lines showing anticancer effects of pharmacological (millimolar) ascorbate, resulting in animal and human trials,⁷ we have found recently that the supplementation of cell culture medium with as little as 5 μM of iron (to reach the minimal level in interstitial fluid) annihilates any anticancer effects of ascorbate. A similar outcome has been observed when the level of iron was corrected via increased percentage of FCS.⁸ Briefly, the combination of iron, ascorbate, and O₂ in the medium generates H₂O₂, which enters and kills cancer cells.⁹ But when iron is present at physiological concentrations it reacts with H₂O₂ to produce hydroxyl radical, which has an extremely low diffusion radius and cannot affect the cells.^{8,10} For a more detailed explanation, we need to take a closer look at the ascorbate–iron–oxygen system, which is composed of two branches (Fig. 1). With ascorbate present in excess (μM), O₂ and iron concentration are the rate limiting factors. High pO_2 in cell culture media promotes branch 1,¹¹ resulting in the accumulation of H₂O₂. However, branch 1 also consumes O₂, so there is a negative feedback present. With the decrease of O₂ concentration, branch 1 slows down and branch

Correspondence to: Ivan Spasojević, Department of Life Sciences, Institute for Multidisciplinary Research, University of Belgrade, Belgrade, Kneza Višeslava 1, Serbia. E-mail: redoxsci@gmail.com




Figure 1 Redox system composed of iron, ascorbate, and molecular oxygen. Two main branches are marked with 1 and 2. Asc, ascorbate; HO•, hydroxyl radical; *Asc, ascorbyl radical.

2 takes over to reduce Fe^{3+} to Fe^{2+} , which further removes H_2O_2 via Fenton chemistry.⁸ Ascorbyl radical is generated in branch 2 and via hydroxyl radical scavenging. The consumption of O_2 and the production of ascorbyl radical are promoted with increasing iron concentrations.⁸ The balance between accumulation and degradation of H_2O_2 is defined by the level of iron, and unfortunately it appears that the major switch between these two is placed between the concentrations of iron in cell culture media and in interstitial fluid. Even more, at *in vivo* O_2 level branch 2 should clearly prevail, not allowing the generation/accumulation of H_2O_2 .

An immediate cause for this commentary is the recently published paper by Du *et al.*, entitled ‘The role of labile iron in the toxicity of pharmacological ascorbate’.¹² The authors have cited our work and recognized that extracellular iron can save cancer cells from ascorbate-related H_2O_2 production but the study, like many before, was conducted in DMEM with 10% FCS. The focus was on the modulation of intracellular level of iron, and it has been presumed that the influx of H_2O_2 from the ascorbate-supplemented medium is (patho)physiologically relevant. The discussion is closed by the following suggestion:

‘However, increasing the level of extracellular as well as intracellular catalytically active labile iron in tumor tissue may enhance the effectiveness of pharmacological ascorbate as an adjuvant in cancer therapy’, although the level of extracellular iron was not taken into consideration as an experimental parameter, and in spite of our findings that the increase of extracellular iron in the medium (from 5 to 10–30 μM ; in fact, we went up to 100 μM – unpublished data) completely prevents anticancer effects of ascorbate.⁸

If we, scientists directly involved in the redox field, continue to keep our eyes closed to this and other technical but fundamental problems, the road is paved for researchers in lucrative fields, such as food industry, pharmacy, and cosmetic industry to develop ‘powerful new antioxidants’, and to compromise our efforts. We cannot act surprised by the fact that promising results from *in vitro* redox studies are rarely translated into success in human clinical trials. For example, at least some phenolic compounds produce H_2O_2 in cell culture media and exert *in vitro* anticancer effects via mechanisms that appear to be similar to ascorbate.^{13,14} So it is no wonder that clear benefits for cancer patients (such as tumor regression or prolonged survival) from the application of (poly) phenolics are still missing in spite of a number of finished clinical trials.¹⁵

To conclude, we urgently need to determine redox properties of interstitial fluid, including the redox activity of iron and the level of endogenous antioxidants and antioxidative enzymes, and to define medium formula and culture maintenance settings that are optimal for redox research. Similarly to medical associations and their guidelines for patient treatment, the societies interested in free radicals might provide recommendations for redox research. Our field has yet much to offer and must not be jeopardized.

Acknowledgement

This study was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, grant number OI173014.

Disclaimer statements

Contributors None.

Funding None.

Conflicts of interest None.

Ethics approval None.

References

- 1 Ganz T, Nemeth E. Hcpidin and disorders of iron metabolism. *Annu Rev Med* 2011;62:347–60.
- 2 Cavill I, Staddon G, Jacobs A. Iron kinetics in the skin of patients with iron overload. *J Invest Dermatol* 1972;58(2):96–8.

- 3 Kakuta K, Orino K, Yamamoto S, Watanabe K. High levels of ferritin and its iron in fetal bovine serum. *Comp Biochem Physiol A Physiol* 1997;118(1):165–69.
- 4 Wesselius LJ, Williams WL, Bailey K, Vamos S, O'Brien-Ladner AR, Wiegmann T. Iron uptake promotes hyperoxic injury to alveolar macrophages. *Am J Respir Crit Care Med* 1999;159(1):100–6.
- 5 Halliwell B. Oxidative stress in cell culture: an under-appreciated problem? *FEBS Lett* 2003;540(1–3):3–6.
- 6 Halliwell B. Cell culture, oxidative stress, and antioxidants: avoiding pitfalls. *Biomed J* 2014;37(3):99–105.
- 7 Parrow NL, Leshin JA, Levine M. Parenteral ascorbate as a cancer therapeutic: a reassessment based on pharmacokinetics. *Antioxid Redox Sign* 2013;19(17):2141–56.
- 8 Mojić M, Bogdanović Pristov J, Maksimović-Ivanić D, Jones DR, Stanić M, Mijatović S, Spasojević I. Extracellular iron diminishes anticancer effects of vitamin C: an *in vitro* study. *Sci Rep* 2014;4:5955.
- 9 Chen Q, Espey MG, Krishna MC, Mitchell JB, Corpe CP, Buettner GR, *et al*. Pharmacologic ascorbic acid concentrations selectively kill cancer cells: action as a pro-drug to deliver hydrogen peroxide to tissues. *Proc Natl Acad Sci USA* 2005;102(38):13604–9.
- 10 Hempel SL, Buettner GR, Wessels DA, Galvan GM, O'Malley YQ. Extracellular iron (II) can protect cells from hydrogen peroxide. *Arch Biochem Biophys* 1996;330(2):401–8.
- 11 Taqui Khan MM, Martell AE. Metal ion and metal chelate catalyzed oxidation of ascorbic acid by molecular oxygen. I. Cupric and ferric ion catalyzed oxidation. *J Am Chem Soc* 1967;89(16):4176–85.
- 12 Du J, Wagner BA, Buettner GR, Cullen JJ. The role of labile iron in the toxicity of pharmacological ascorbate. *Free Radic Biol Med* 2015;88:289–95.
- 13 Long LH, Clement MV, Halliwell B. Artifacts in cell culture: rapid generation of hydrogen peroxide on addition of (–)-epigallocatechin, (–)-epigallocatechin gallate, (+)-catechin, and quercetin to commonly used cell culture media. *Biochem Biophys Res Commun* 2000;273(1):50–3.
- 14 Long LH, Kirkland D, Whitwell J, Halliwell B. Different cytotoxic and clastogenic effects of epigallocatechin gallate in various cell-culture media due to variable rates of its oxidation in the culture medium. *Mutat Res* 2007;634(1–2):177–83.
- 15 Carocho M, Ferreira IC. The role of phenolic compounds in the fight against cancer – a review. *Anticancer Agents Med Chem* 2013;13(8):1236–58.